

<b>Subject:</b>	Epithelial Cell Cytology in Breast Cancer Risk Assessment		
<b>Policy Number:</b>	PO-RE-032v4		
<b>Effective Date:</b>	02/01/2026	<b>Last Approval Date:</b>	12/15/2025

[Policy Description](#) | [Indications and/or Limitations of Coverage](#) | [Scientific Background](#) |  
[Guidelines and Recommendations](#) | [Applicable Codes](#) | [Definitions](#) | [Related Policies](#) |  
[Reference Materials](#) | [Revision History](#) | [Disclaimer](#)

## I. Policy Description

Nipple aspiration and/or ductal lavage are non-invasive techniques to obtain epithelial cells for cytological examination to aid in the evaluation of nipple discharge for breast cancer risk.<sup>1</sup> Fine needle aspiration (FNA) is another approach that can be used in the initial diagnosis of a suspicious breast mass, although core biopsy is superior in sensitivity, specificity, and correct histological grading.<sup>2</sup>

### Indications and/or Limitations of Coverage

Application of coverage criteria is dependent upon an individual's benefit coverage at the time of the request. Specifications pertaining to Medicare and Medicaid can be found in the "Applicable State and Federal Regulations" section of this policy document.

*The following does not meet coverage criteria due to a lack of available published scientific literature confirming that the test(s) is/are required and beneficial for the diagnosis and treatment of an individual's illness.*

1. Cytologic analysis of epithelial cells to assess breast cancer risk and manage patients at high risk of breast cancer **DOES NOT MEET COVERAGE CRITERIA.**

### Scientific Background

Breast cancer is the most frequently diagnosed cancer and is a leading cause of cancer death in the United States. Nipple discharge is a common breast complaint. Most nipple discharge is of benign origin; however, it is necessary to differentiate patients with benign nipple discharge from those who have an underlying pathology. In approximately five to 20 percent of pathologic nipple discharge cases, cancer is identified.<sup>1</sup>

Breast cancer originates in breast epithelium and is associated with progressive molecular and morphologic changes. Individuals with atypical breast ductal epithelial cells have an increased relative

risk of breast cancer. Cytological evaluation of epithelial cells in nipple discharge has been used as a diagnostic aid. Due to the scant cellularity of specimens by expression or aspiration of nipple discharge, ductal lavage was developed to enhance the ease and efficiency of collecting breast epithelial cells for cytologic analysis. The analysis of breast-specific liquid biopsies, such as nipple aspirate fluid, has potential to be used as a biomarker profiling technique for monitoring breast health.<sup>3</sup> Researchers report that the measurement of nipple aspirate fluid, including miRNA, pathological nipple discharge, and breast ductal fluids, may help to improve early detection and management of breast cancer.<sup>4</sup>

Fine needle aspiration (FNA) is a biopsy option for a suspicious palpable breast mass. FNA is a rapid diagnosis technique, but it is not as accurate as core needle biopsy. FNA cannot differentiate in situ and invasive cancer and has higher rates of negative results and insufficient samples than core needle biopsy. The success of FNA results also varies with the operator and cytopathologist.<sup>5</sup>

### ***Analytic Validity***

In a retrospective study of 618 patients with nipple discharge over a 14-year period, the sensitivity and specificity of cytology were 17 and 66 percent, respectively; the authors concluded that "nipple discharge cytology has little complementary diagnostic value."<sup>6</sup>

### ***Clinical Utility and Validity***

Hornberger, et al. (2015) performed a meta-analysis on the use of nipple aspirate fluid (NAF) in identifying breast cancer based on proliferative epithelial disease (PED). The authors reviewed 16 articles, 20808 unique aspirations, and 17378 subjects. Among cancer-free patients, 51.5% aspirations contained fluid, of which 27.7% showed a PED on cytology. Of the two prospective studies of 7850 women, patients with abnormal cytology showed a 2.1-fold higher risk of developing breast cancer compared to those without fluid.<sup>7</sup>

Chatterton, et al. (2016) measured sex steroid levels in NAF; hormones were measured in samples from 160 breast cancer cases and 157 controls. Results showed a significantly higher concentration of dehydroepiandrosterone (DHEA) in the NAF of patients with breast cancer compared to controls; further, DHEA levels were highly correlated with estradiol levels, indicating "a potentially important role of this steroid in breast cancer risk."<sup>8</sup>

Kamalı and Kamalı (2022) studied the usefulness of testing methods in surgical decision making. The study included 141 patients with pathological nipple discharge who were planning to undergo surgery. The diagnostic efficiency of ductal lavage cytology was compared to that of ultrasonography, mammography, magnetic resonance imaging, and ductography. The sensitivity of ductal lavage cytology was 70.5% and the specificity was 94.1%. The authors conclude that "negative cytology does not exclude the possibility of malignancy, and positive results do not help in the differential diagnosis."<sup>9</sup>

Zhu, et al. (2025) released a study on using direct analysis of nipple discharge using liquid based cytology (LBC) and by passing the conventional smear cytology (CSC). "LBC has better diagnostic

sensitivity than CSC (40.00% vs. 22.22%,  $\chi^2 = 6.636$ ,  $P = 0.01$ ). The specificity was improved (LBC 100% vs. CSC 95.2%)." When looking at the diagnostic validity "LBC had higher sensitivity than CSC for breast tumors (52.78% vs. 30.56% for breast cancer; and 31.48% vs. 16.67% for intraductal papilloma)." <sup>10</sup> The study concluded that LBC may overcome the contamination and low-cellularity limitations associated with smear cytology and improve the diagnostic yield in patients presenting with nipple discharge.

## Guidelines and Recommendations

### American Society of Breast Surgeons (ASBS)

The Official Statement by the ASBS regarding Screening Mammography does not mention ductal lavage at all in their statement.<sup>11</sup>

In 2016, the ASBS published a consensus guideline on the concordance assessment of image-guided breast biopsies and the management of borderline or high-risk lesions. These guideline state that "the decision to excise a papillary lesion without atypia needs to be individualized based on risk, including such criteria as size; symptomatology, including palpability and presence of nipple discharge; and breast cancer risk factors."<sup>12</sup> This is the only mention of nipple discharge in the document.

### National Comprehensive Cancer Network (NCCN)

National Comprehensive Cancer Network Clinical Practice Guidelines in breast cancer screening and diagnosis state that "thermography and ductal lavage are not recommended by the NCCN Panel for breast cancer screening or diagnosis."<sup>13</sup> The NCCN also notes that "the FDA has issued a safety alert stating that ductal lavage should not be a replacement for mammograms."<sup>13</sup>

### Food and Drug Administration (FDA)

In 2017 the FDA issued a safety warning stating that "...the FDA is unaware of any valid scientific data to show that a nipple aspirate test, when used on its own, is an effective screening tool for any medical condition, including the detection of breast cancer or other breast disease."<sup>14</sup> This was further affirmed with a safety warning published in 2023: "thermograms and nipple aspirate tests are not substitutes for mammograms."<sup>15</sup>

### American College of Radiology (ACR)

The 2022 ACR appropriateness criteria for the evaluation of nipple discharge do not mention cytology. The ACR states that "image-guided FNA and core biopsy are not required for the evaluation of physiologic nipple discharge" but "image-guided FNA and core biopsy are not required for the evaluation of physiologic nipple discharge." The ACR also notes "although some institutions demonstrate good results using FNA, larger series have shown that core biopsy is superior to FNA in terms of sensitivity, specificity, and correct histologic grading of a lesion."<sup>16</sup>

### National Cancer Institute (NCI)

The NCI notes that FNA, nipple aspirate, and ductal lavage are proposed as screening techniques for breast cancer “but none have demonstrated an association with reduced mortality.”<sup>17</sup> These methods are not recommended as routine screening tools and are not included in NCI’s endorsed clinical practices.

### Food and Drug Administration (FDA)

Many labs have developed specific tests that they must validate and perform in house. These laboratory-developed tests (LDTs) are regulated by the Centers for Medicare and Medicaid (CMS) as high-complexity tests under the Clinical Laboratory Improvement Amendments of 1988 (CLIA '88). LDTs are not approved or cleared by the U. S. Food and Drug Administration; however, FDA clearance or approval is not currently required for clinical use.

## II. Applicable Codes

Code	Description	Comment
88108	Cytopathology, concentration technique, smears and interpretation (e.g., Saccomanno technique)	
88112	Cytopathology, selective cellular enhancement technique with interpretation (e.g., liquid based slide preparation method), except cervical or vaginal	
88172	Cytopathology, evaluation of fine needle aspirate; immediate cytohistologic study to determine adequacy for diagnosis, first evaluation episode, each site	
88173	Cytopathology, evaluation of fine needle aspirate; interpretation and report	
88177	Cytopathology, evaluation of fine needle aspirate; immediate cytohistologic study to determine adequacy for diagnosis, each separate additional evaluation episode, same site (List separately in addition to code for primary procedure)	

## III. Definitions

Term	Meaning
N/A	N/A


## IV. Related Policies

Policy Number	Policy Description
PO-RE-041	Serum Tumor Markers for Malignancies
PO-RE-032	Use of Common Genetic Variants (Single Nucleotide Polymorphisms) To Predict Risk of Non-Familial Breast Cancer

Current Procedural Terminology © American Medical Association. All rights reserved.

*Procedure codes appearing in Reimbursement Policy documents are included only as a general reference tool for each policy. They may not be all-inclusive.*

## V. Reference Materials

Golshan M. Nipple discharge. Updated February 12, 2024. <a href="https://www.uptodate.com/contents/nipple-discharge">https://www.uptodate.com/contents/nipple-discharge</a>
Moy L, Heller SL, Bailey L, et al. ACR Appropriateness Criteria® Palpable Breast Masses. J Am Coll Radiol. May 2017;14(5s):S203-s224. doi:10.1016/j.jacr.2017.02.033
Shaheed SU, Tait C, Kyriacou K, Linforth R, Salhab M, Sutton C. Evaluation of nipple aspirate fluid as a diagnostic tool for early detection of breast cancer. Clin Proteomics. 2018;15:3. doi:10.1186/s12014-017-9179-4
Moelans CB, Patuleia SIS, van Gils CH, van der Wall E, van Diest PJ. Application of Nipple Aspirate Fluid miRNA Profiles for Early Breast Cancer Detection and Management. Int J Mol Sci. Nov 19 2019;20(22)doi:10.3390/ijms20225814
Joe BN, Esserman LJ. Breast biopsy. Updated May 3, 2023. <a href="https://www.uptodate.com/contents/breast-biopsy">https://www.uptodate.com/contents/breast-biopsy</a>
Kooistra BW, Wauters C, van de Ven S, Strobbe L. The diagnostic value of nipple discharge cytology in 618 consecutive patients. European journal of surgical oncology : the journal of the European Society of Surgical Oncology and the British Association of Surgical Oncology. Jun 2009;35(6):573-7. doi:10.1016/j.ejso.2008.09.009
Hornberger J, Chen SC, Li Q, Kakad P, Quay SC. Proliferative epithelial disease identified in nipple aspirate fluid and risk of developing breast cancer: a systematic review. Current medical research and opinion. Feb 2015;31(2):253-62. doi:10.1185/03007995.2014.988209
Chatterton RT, Heinz RE, Fought AJ, et al. Nipple Aspirate Fluid Hormone Concentrations and Breast Cancer Risk. Horm Cancer. Apr 2016;7(2):127-36. doi:10.1007/s12672-016-0252-7
Kamalı GH, Kamalı S. The Role of Ductal Lavage Cytology in the Diagnosis of Breast Cancer.

Archives of Iranian Medicine (AIM). 2022;25(11)doi:10.34172/aim.2022.118
Zhu J, Cong H, Zhang X, et al. New method for cytological evaluation using direct nipple discharge without aspiration. <i>Scientific Reports</i> . 2025/02/04 2025;15(1):4175. doi:10.1038/s41598-025-88456-9
ASBS. Screening Mammography. 2019. <a href="https://www.breastsurgeons.org/docs/statements/Position-Statement-on-Screening-Mammography.pdf">https://www.breastsurgeons.org/docs/statements/Position-Statement-on-Screening-Mammography.pdf</a>
ASBS. Consensus Guideline on Concordance Assessment of Image-Guided Breast Biopsies and Management of Borderline or High-Risk Lesions. <a href="https://www.breastsurgeons.org/docs/statements/Consensus-Guideline-on-Concordance-Assessment-of-Image-Guided-Breast-Biopsies.pdf">https://www.breastsurgeons.org/docs/statements/Consensus-Guideline-on-Concordance-Assessment-of-Image-Guided-Breast-Biopsies.pdf</a>
NCCN. NCCN Clinical Practice Guidelines in Oncology; Breast Cancer Screening and Diagnosis V2.2025. National Comprehensive Cancer Network. Updated March 28, 2025. <a href="https://www.nccn.org/professionals/physician_gls/pdf/breast-screening.pdf">https://www.nccn.org/professionals/physician_gls/pdf/breast-screening.pdf</a>
Williams A, Omnipographics Inc. Breast cancer sourcebook : basic consumer health information about the prevalence, risk factors, and symptoms of breast cancer, including ductal and lobular carcinoma in situ, invasive carcinoma, inflammatory breast cancer, and breast cancer in men and pregnant women; along with facts about benign breast changes, breast cancer screening and diagnostic tests, treatments such as surgery, radiation therapy, chemotherapy, and hormonal and biologic therapies, tips on managing treatment side effects and complications, a glossary of terms, and a directory of resources for additional help and information. Sixth edition. ed. Omnipographics,; 2019:1 online resource.
FDA. Mammography: What You Need to Know. Updated October 26, 2023. <a href="https://www.fda.gov/consumers/consumer-updates/mammography-what-you-need-know">https://www.fda.gov/consumers/consumer-updates/mammography-what-you-need-know</a>
Sanford MF, Slanetz PJ, Lewin AA, et al. ACR Appropriateness Criteria® Evaluation of Nipple Discharge: 2022 Update. <i>J Am Coll Radiol</i> . Nov 2022;19(11s):S304-s318. doi:10.1016/j.jacr.2022.09.020
NCI. Breast Cancer Screening (PDQ®)–Health Professional Version. Updated April 10, 2025. <a href="https://www.cancer.gov/types/breast/hp/breast-screening-pdq">https://www.cancer.gov/types/breast/hp/breast-screening-pdq</a>

## VI. Revision History

Revision Date	Summary of Changes
09/04/2025	Reviewed and Updated: Updated the background, guidelines and recommendations, and evidence-based scientific references. Literature review did not necessitate any modifications to coverage criteria.
09/04/2024	Reviewed and Updated: Updated the background, guidelines and recommendations, and evidence-based scientific references. Literature review did not necessitate any modifications to coverage criteria.
09/06/2023	Reviewed and Updated: Updated the background, guidelines and recommendations, and evidence-based scientific references. Literature review necessitated the following changes to coverage criteria: CC1 updated to reflect that all cytological analysis for breast cancer diagnosis DNMCC, as biopsy should be used to diagnose. CC now reads: "1) Cytologic

	analysis of epithelial cells to assess breast cancer risk and manage patients at high risk of breast cancer DOES NOT MEET COVERAGE CRITERIA.” Added CPT code 88172, 88173, 88177
--	---

## Disclaimer

Healthfirst's claim edits follow national industry standards aligned with CMS standards that include, but are not limited to, the National Correct Coding Initiative (NCCI), the National and Local Coverage Determination (NCD/LCD) policies, appropriate modifier usage, global surgery and multiple procedure reduction rules, medically unlikely edits, duplicates, etc. In addition, Healthfirst's coding edits incorporate industry-accepted AMA and CMS CPT, HCPCS and ICD-10 coding principles, National Uniform Billing Editor's revenue coding guidelines, CPT Assistant guidelines, New York State-specific coding, billing, and payment policies, as well as national physician specialty academy guidelines (coding and clinical). Failure to follow proper coding, billing, and/or reimbursement policy guidelines could result in the denial and/or recoupment of the claim payment.

This policy is intended to serve as a resource for providers to use in understanding reimbursement guidelines for professional and institutional claims. This information is accurate and current as of the date of publication. It provides information from industry sources about proper coding practice. However, this document does not represent or guarantee that Healthfirst will cover and/or pay for services outlined. Reimbursement decisions are based on the terms of the applicable evidence of coverage, state and federal requirements or mandates, and the provider's participation agreement. This includes the determination of any amounts that Healthfirst or the member owes the provider.