

# Documentation and Coding: Heart Failure

Created October 2021

At Healthfirst, we are committed to helping providers accurately document and code their patients' health records. This tip sheet is intended to assist providers and coding staff with the documentation and ICD-10-CM selection on services submitted to Healthfirst specifically for coding heart failure. It provides information from industry sources about proper coding practice. However, this document does not represent or guarantee that Healthfirst will cover and/or pay for services outlined. Coverage decisions are based on the terms of the applicable evidence of coverage and the provider's participation agreement. This includes the determination of any amounts that Healthfirst or the member owes the provider.

Type of Heart Failure					Clinical Documentation Should Include	
Left Ventricular Failure, Unspecified I50.1	Systolic (Congestive) I50.2*	Diastolic (Congestive) I50.3*	Combined Systolic (Congestive) & Diastolic (Congestive) I50.4*	Other/Right Heart Failure I50.81*	<b>Status of Condition</b>	Stable; Improved; Worsening; Compensated Exacerbation
					<b>Severity</b>	Systolic; Diastolic; Combined
<b>Heart Failure with Other Chronic Conditions</b>					<b>Type of Heart Failure</b>	Acute; Chronic; Acute-on-chronic
<b>Hypertensive Heart Disease</b>	<b>Hypertensive Heart &amp; Chronic Kidney Disease (CKD)</b>			<b>Rheumatic Heart Failure</b>	<b>Link Associated Conditions/Manifestations</b>	"Due to" "Secondary to" "Associated with"
With Heart Failure I11.0 Without Heart Failure I11.9	Without Heart Failure, with CKD Stage 1-4 or Unspecified CKD - I13.10 Without Heart Failure, with CKD Stage 5 or End-Stage Renal Disease (ESRD) - I13.11 With Heart Failure, with Stage 5 CKD or ESRD - I13.12			Rheumatic heart failure I09.81	<b>Interpretation of Diagnostic Tests</b>	Catheterization Cardiac stress testing Echocardiogram, ECG or EKG, X-ray, CT/MRI scans or Nuclear heart scans
<b>Other Types of Heart Failure</b>					<b>Any risk factors</b>	Smoking, Obesity, Congenital Heart Disease, Abnormal Heart Valves or Diseases of Heart Muscle, Past Heart Attack
<ul style="list-style-type: none"> <li>• Right heart failure due to left heart failure - I50.814, code also the type of left ventricular failure, from I50.2–I50.43 if known.</li> <li>• Patients with biventricular failure can have right heart disease due to one cause and left heart disease due to another.               <ul style="list-style-type: none"> <li>- Biventricular heart failure - I50.82, code when there is a different disease-causing heart failure in each ventricle, also assign the type of left ventricular failure I50.2–I50.43 if known.</li> <li>- High output heart failure - I50.83, occurs when the high demand for blood exceeds the capacity of a normally functioning heart to meet the demand.</li> <li>- End stage heart failure (Stage D of heart failure) - I50.84, code also type of heart failure as systolic or diastolic, I50.2-I50.43 if known.</li> </ul> </li> <li>• Code first the chronic conditions and assign additional code from category I50 to identify type of heart failure. Add an additional code from category N18 to identify the stage of CKD.</li> </ul>					<b>Procedure/ Postprocedural of Heart Failure</b>	i.e., Post cardiectomy syndrome; Postmastectomy lymphedema syndrome; Postprocedural hypertension; Type of surgery - following heart catheterization
					<b>Treatment plans</b>	Link medications to condition, Specialty Referrals, Consultations Requested, Device Therapy Treatment for the cause of heart failure

\*Requires an additional digit to complete the diagnosis code.

# Documentation and Coding: Heart Failure

## Documentation Tips

- Heart failure and congestive heart failure (CHF) classify to the same ICD-10-CM I50\* category.
- When heart failure is described as decompensated or exacerbated, it should be coded as **acute-on-chronic**.
- Document heart failure to the highest level of specificity, i.e., congestive, hypertensive, post-operative, acute, chronic, acute-on-chronic, diastolic, systolic, etc.
- Ensure that the results of an echocardiogram differentiating between systolic and diastolic heart failure is documented for appropriate diagnosis code selection.
- When the final diagnosis is CHF along with either systolic or diastolic heart failure, **only code the type of heart failure (systolic and/or diastolic)**.
- Code assignment for CHF is dependent upon **both the type of failure (e.g., left systolic, diastolic, combined) and severity (e.g., acute chronic, acute on chronic)**. If not present, the physician should be queried. Code I50.9 should only be reported if the type of heart failure cannot be further specified.
- ICD-10 classification presumes a causal relationship between hypertension and heart involvement and between hypertension and kidney involvement, as these conditions should be coded as related unless the documentation clearly states the conditions are unrelated (Guideline I.C.9.a).

\*Requires an additional digit to complete the diagnosis code.

## Coding Examples:

Documentation	Patient is here for a follow-up of chronic systolic heart failure. HPI: No chest pain, dyspnea on exertion, denies dizziness, non-smoker. PE: NAD no JVD Carotid 2+, bruits. Heart s1 s2 irregular, no gallop, chest is clear. Extremities no edema femoral pulses 2+. Diagnostic studies EKG: Atrial fib poor R wave progression vi3 nonspecific ST-T abnormalities. A/P: chronic systolic CHF – compensated. Atrial fibrillations controlled ventricular response.
Diagnosis reported	I50.22 - Chronic systolic (congestive) heart failure I48.91 - Unspecified atrial fibrillation
Rationale	The diagnosis codes are supported with MEAT (Monitored, Evaluated, Assessed or Treated (MEAT)).

## Documentation and Coding: Heart Failure

### Coding Examples: *(continued)*

<b>Documentation</b>	The patient, who has a history of heart failure, hypertension, and coronary artery disease, presents with ST-elevation myocardial infarction. As a secondary diagnosis, the provider recorded HFmrEF, meaning heart failure with mid-range or mildly reduced ejection fraction (EF). How should heart failure with mid-range or mildly reduced EF be coded?
<b>Diagnosis reported</b>	I50.22 - Chronic systolic (congestive) heart failure
<b>AHA Coding Clinic 3rd QTR 2020 Volume 7 Rationale</b>	The ejection fraction indicates the amount of blood that is pumped out from the ventricle to the body during systole (the phase in which the heart muscle contracts). When a patient with a history of heart failure is described in terms of reduced ejection fraction (mid-range or mildly), assign a code for chronic systolic heart failure (I50.22).

<b>Documentation</b>	Follow-up visit for Hypertensive heart w/chronic diastolic CHF and CKD4. HPI: No chest pain, dyspnea on exertion, non-smoker. PE: NAD no JVD Carotid 2+, bruits. Heart s1 s2 irregular, no gallop, chest is clear. Extremities 2+ edema; GU no urinary retention; A/P: Hypertensive heart disease with chronic diastolic CHF, CKD stage 4.
<b>Diagnosis reported</b>	I13.0 - Hypertensive heart and CKD with heart failure and stage 1 through 4 CKD, or unspecified CKD. I50.32 - Chronic diastolic (congestive) heart failure N18.4 - Chronic kidney disease, stage 4 (severe)
<b>Rationale</b>	ICD-10 Guidelines assign codes from combination category I13, Hypertensive heart and CKD, when there is hypertension with both heart and kidney involvement. If heart failure is present, assign an additional code from category I50 to identify the type of heart failure. The appropriate code from category N18, CKD, should be used as a secondary code with a code from category I13 to identify the stage of CKD.

# Documentation and Coding: Heart Failure

---

## Common coding practices that providers should avoid:

- Do not document heart failure as a confirmed condition if it is suspected. Instead, document signs and symptoms in the absence of a confirmed diagnosis.
- Do not describe heart failure as “history of” if the condition is still active (in diagnosis “history of” implies a condition that no longer exists as a current problem).
- Do not use words that imply uncertainty (“likely,” “probable,” “apparently,” “consistent with,” etc.) to describe a current or confirmed diagnosis.
- Do not document temporary or transient heart failure that occurred in the past and is no longer present as current.

## Questions?

Contact us at [#Risk\\_Adjustments\\_and\\_clinical\\_Documentation@healthfirst.org](mailto:Risk_Adjustments_and_clinical_Documentation@healthfirst.org).

For additional documentation and coding guidance, please visit the Coding section at [hfproviders.org](http://hfproviders.org).

### References:

[AHIMA.org](http://AHIMA.org); [CodingClinicAdvisor.com](http://CodingClinicAdvisor.com); [ICD-10-CM-Guidelines2021](#)